



3112 Fairview Drive
Owensboro, KY 42303
Phone: 270-922-2500
Fax: 270-922-2505
info@Landmarkgi.com

Miguel Lalama, MD, MPH

Brittany Davidson, APRN

Simone McCormack, APRN

ERCP

Patient: _____

Procedure Date: _____

Location of Procedure:

Ohio County Hospital

1211 Old Main St, Hartford

Owensboro Surgery Center

1000 Breckenridge St, Suite 100

Owensboro Health Reg. Hospital

1201 Pleasant Valley Rd

Arrival Time: _____

Your physician has recommended that you have an EGD. This is an exam for your upper GI tract. Your physician will also perform an ERCP which uses a dye to highlight the bile ducts on X-ray images. **Please read the following directions prior to your procedure date. If you have any questions or concerns regarding the instructions or prep directions, please contact the office prior to your procedure.**

SURGERY PREP INSTRUCTIONS

FOR FIVE (5) DAYS BEFORE YOUR PROCEDURE:

- Do NOT take products that contain iron -- such as multivitamins. Other common names for iron are ferrous sulfate, ferrous gluconate, ferrous fumarate, Femiron, or Niferex.
- Stop taking fiber supplements. This includes Metamucil, Fibercon, Citrucel, Konsyl, Benefiber, Fibersure, and Colon Care Capsules.
- DO NOT EAT high-fiber foods like nuts, seeds, popcorn, beans, peas, raw salad, celery, tomatoes.
- DO NOT EAT wheat germ, granola, cereal with visible grains/fruit, bread with seeds/ visible grains.
- YOU MAY eat cooked, low-fiber vegetables like carrots, broccoli, cauliflower, potatoes with no skin.
- YOU MAY eat fruit, but skin and seeds need to be removed before eating.

THE DAY BEFORE YOUR PROCEDURE:

- Except for the above foods, you may eat like normal the day before your procedure until MIDNIGHT. For your safety and the accuracy of these procedures, DO NOT EAT AFTER midnight. This includes gum, mints, water, alcohol, and coffee.

THE DAY OF YOUR PROCEDURE:

- You should take your cardiac, blood pressure, and/or seizure medications with a sip of water.
- DO NOT EAT OR DRINK anything on the day of your procedure.

Sedation:

You will receive sedation for your procedure. This helps you relax and feel comfortable during the procedure. When you have sedation, you must follow the instructions listed below or your procedure will be delayed or canceled. You may feel fine after your procedure, but your judgment may be impaired from the sedation medication you receive.

- Arrange to have an adult family member or friend accompany you to your procedure and to other appointments, pick you up from the procedure unit, and drive you home or to your place of lodging.
- The absence of a responsible adult will result in a delay or cancellation of your procedure if plans have not been made for your after-procedure travel.
- Do not drive for the rest of the day. Unaccompanied air travel on the same day as your procedure is not permitted. This is for your safety and the safety of others.

Medications:

If you have Diabetes: Due to fasting prior to the procedure, if you are insulin dependent, you will take **HALF** of your normal morning dose the morning of the procedure. For further questions or concerns regarding insulin, please contact the ordering provider.

If you take blood thinning medications: Blood-thinning medications affect clotting and bleeding. Both the health care provider who manages these medications and the provider doing your procedure will need to decide if your medications need to change. Please make sure that the provider is aware if you are on any of these medications and make sure that you receive directions on when to stop taking these prior to the procedure. Some medications need to be stopped up to 7 days prior to the procedure, so make sure that you have plenty of time to receive these directions. **DO NOT STOP THESE MEDICATIONS WITHOUT INSTRUCTIONS FROM A PROVIDER.**

If you take daily medication(s) outside of the medications listed above, you may do so up until the day prior to the procedure. The morning of your procedure you may take any cardiac (heart), including blood pressure medications, and/or seizure medications with a small sip of water.

If you are taking any weight loss medications: Please notify the provider if you are taking any of the following medications: phentermine (Adipex-P), lorcaserin (Belviq), phendimetrazine (Bontril). You will be provided instructions on how long to hold these prior to your procedure.